



The Role of Macrobenthic Indicator Species as Bioindicators for Assessing Marine Ecosystem Health

Ali Abbasi^{a*}, Mehran Loghmani^a

^aDepartment of Marine Biology, Faculty of Marine Sciences, Chabahar University of Maritime & Marine Sciences, Chabahar, Sistan & Baluchestan, Iran.

*Corresponding author: aliabbasi1376.2014@gmail.com, <https://orcid.org/0009-0005-4934-0158>

Article Info

ABSTRACT

Article type:
Research Article

Article history:
Received: 2026/01/27
Received in revised form:
2026/02/19
Accepted 2026/02/29
Available online: 2026/03/06

Keywords:
Marine ecosystem,
Marine pollution,
Environmental monitoring,
Biodiversity,
Macrobenthic indicator
species.

Marine ecosystems are under increasing pressure from human activities & climate change, making effective monitoring more important than ever. Among the available tools, macrobenthic indicator species stand out as practical & reliable options for assessing environmental conditions in coastal & offshore waters. This review looks at what these organisms are, why they matter ecologically, & how they can be used in real-world monitoring programs. We cover the main taxonomic groups involved, including mollusks, crustaceans, annelids, echinoderms, & demersal fish, & discuss their specific roles in environmental assessment. Macrobenthic organisms work well as bioindicators for several reasons. They don't move around much, they live long enough to show cumulative effects, they respond clearly to environmental changes, & they build up pollutants in their tissues over time. These traits make them useful for judging habitat quality, tracking ecosystem health, & spotting potential problems before they get worse. Methods like IndVal, AMBI, & M-AMBI have helped make benthic assessments more reliable by adding quantitative rigor to what used to be mostly descriptive work. The literature shows that different groups serve different purposes. Polychaetes like *Capitella capitata* & *Heteromastus filiformis* are well known as signs of organic enrichment & sediment contamination. Filter-feeding bivalves are good for long-term heavy metal monitoring because they concentrate contaminants in their tissues. Amphipods respond quickly to sediment toxicity, while echinoderm larvae are sensitive to various stressors. Demersal fish add another layer by accumulating pollutants at higher positions in the food web. What makes this review different is that it brings together ecological perspectives on macrobenthic indicators while also pointing out how newer biotechnological tools could improve species identification & monitoring accuracy. Putting together regional databases of these species would make future work easier & help build more effective conservation strategies. The review also highlights that while these methods are widely used, there's still room for improvement in how species are selected & how results are interpreted across different regions.

Introduction

Marine ecosystems are increasingly exposed to a wide range of anthropogenic pressures, including coastal development, industrial pollution, eutrophication, overfishing, habitat degradation, & climate change. These stressors have altered the structure & functioning of benthic habitats & have intensified the need for reliable environmental monitoring tools. Coastal & offshore marine environments provide essential ecosystem services, including nutrient cycling, carbon sequestration, food provision, & shoreline protection, all of which are critical for human well-being & economic development. However, the growing intensity & frequency of human activities in marine environments have led to unprecedented rates of biodiversity loss, habitat fragmentation, & ecosystem dysfunction, prompting urgent calls for effective monitoring & management strategies.

Among the available biological indicators, macrobenthic organisms have received considerable attention due to their ecological importance, relatively sedentary lifestyle, broad distribution, & high sensitivity to environmental disturbances. Benthic communities inhabit the seafloor & are directly exposed to contaminants, organic enrichment, & physical disturbances that accumulate in sediments. Their limited mobility prevents them from avoiding unfavorable conditions, making them reliable indicators of local environmental quality. Additionally, macrobenthic organisms exhibit a wide range of life-history traits, feeding strategies, & physiological tolerances, allowing them to reflect ecological conditions across different spatial & temporal scales. Benthic communities are also relatively easy to sample using standard grab or corer techniques, & their taxonomic identification is well-established for most major groups, facilitating standardized monitoring programs worldwide.

Macrobenthic indicator species can provide valuable information about ecosystem health by reflecting both short-term & long-term ecological changes. The presence, absence, abundance, & diversity of benthic species respond to variations in sediment characteristics, organic matter loading, oxygen availability, & contaminant concentrations. Opportunistic species often proliferate under disturbed conditions, while sensitive species decline or disappear, resulting in predictable shifts in community composition that can be quantified using ecological indices. This capacity to integrate ecological stressors over time makes macrobenthic indicators particularly valuable for assessing cumulative impacts & tracking recovery trajectories following management interventions.

The present review aimed to investigate the concept of macrobenthic indicator species, examine their ecological significance, identify major taxonomic groups used in marine environmental assessment, & evaluate their applications in monitoring marine ecosystem quality. Furthermore, this study highlights the advantages, limitations, & future prospects of using macrobenthic indicators in environmental management & conservation programs.

Method

This study was conducted as a comprehensive review of scientific literature related to macrobenthic indicator species & their applications in marine environmental monitoring. A systematic literature search was performed using major scientific databases, including Google Scholar, Scopus, Web of Science, & ResearchGate, to ensure broad coverage of peer-reviewed publications. The literature search was performed using keywords such as “indicator species,” “macrobenthos,” “marine ecosystem monitoring,” “ecological indicators,” & “benthic community structure.” Boolean operators were used to combine search terms & refine the results. The search was limited to English-language publications, & no specific time restrictions were applied to capture both foundational studies & recent advances. Studies were selected based on scientific quality, relevance to the research topic, publication in peer-reviewed journals, & contribution to ecological assessment & environmental monitoring.

Preference was given to studies that provided quantitative data, validated methodologies, & practical applications in different marine regions.

Information extracted from the selected studies was categorized & analyzed to identify common patterns, methodological approaches, major findings, & existing challenges. Particular attention was given to the ecological responses of macrobenthic communities to environmental stressors, the development of indicator-species methodologies, & the application of ecological indices such as Indicator Value Analysis (IndVal), AZTI Marine Biotic Index (AMBI), & Multivariate AMBI (M-AMBI). The extracted data were synthesized to develop a comprehensive overview of the current state of knowledge, identify methodological strengths & weaknesses, & propose directions for future research. The review also considered the geographic distribution of studies to assess the regional applicability of different indicator approaches & identify areas where knowledge is limited.

Results

The reviewed studies demonstrate that various macrobenthic groups, including polychaetes, mollusks, crustaceans, echinoderms, cnidarians, & demersal fishes, serve as effective indicators of environmental quality in marine ecosystems. Polychaete worms, particularly opportunistic species such as *Capitella capitata* & *Heteromastus filiformis*, are widely recognized as indicators of organic enrichment, sediment contamination, & hypoxic conditions. These species are often among the first to colonize disturbed sediments & can rapidly proliferate under high organic loading, making them sensitive indicators of eutrophication & wastewater discharge. Mollusks, especially filter-feeding bivalves, are capable of accumulating heavy metals & other pollutants, making them valuable biomonitors for long-term contamination assessment. Their sedentary nature & feeding behavior result in direct exposure to sediment-associated contaminants, allowing them to reflect bioavailable pollutant concentrations in the environment. Amphipod crustaceans are sensitive to sediment toxicity & organic pollution, whereas echinoderms exhibit strong responses to environmental stress during their larval stages & can indicate habitat deterioration. Demersal fish also contribute to environmental monitoring through the bioaccumulation of contaminants within higher trophic levels, providing an integrated measure of ecosystem-level contamination.

The findings further reveal that macrobenthic communities are strongly influenced by environmental variables such as sediment grain size, organic matter content, dissolved oxygen concentration, salinity, water depth, & hydrodynamic conditions. The loss of sensitive taxa & the proliferation of tolerant species lead to simplified community structures & reduced functional diversity, which can impair ecosystem functioning & resilience.

The review also highlights the growing application of quantitative methods for identifying indicator species. Approaches such as IndVal combine species abundance & frequency of occurrence to determine species-environment relationships & improve ecological assessments. Similarly, AMBI & M-AMBI have become widely used tools for evaluating ecological status based on benthic community composition. These indices assign species to ecological groups based on their sensitivity to disturbance & enable the classification of sites into quality classes ranging from undisturbed to heavily impacted.

Conclusions

The review confirms that macrobenthic indicators offer a practical & scientifically robust approach to marine environmental monitoring. Their ecological relevance, cost-effectiveness, & ability to integrate environmental conditions over time make them invaluable for detecting & quantifying anthropogenic impacts. However, successful implementation requires careful consideration of indicator selection, methodological standardization, & regional calibration. The development of comprehensive databases & continuous refinement of ecological indices will further strengthen the utility of macrobenthic indicators in environmental management.

Macrobenthic indicator species represent powerful & cost-effective tools for assessing marine environmental quality & ecosystem health. Their sensitivity to environmental changes, limited mobility, long life span, & capacity to integrate ecological conditions over time make them particularly suitable for monitoring anthropogenic impacts & climate-related disturbances. The use of indicator species can facilitate habitat assessment, early detection of environmental degradation, ecological risk evaluation, & the development of conservation strategies. However, improving the reliability of biological assessments requires careful species selection, incorporation of local ecological knowledge, long-term monitoring programs, & the application of advanced analytical & biotechnological techniques. The development of comprehensive regional & global databases for macrobenthic indicator species, together with the refinement of ecological indices, will significantly enhance future marine monitoring efforts & support sustainable management of marine ecosystems in the face of increasing environmental challenges. Future research should focus on integrating molecular & traditional taxonomic approaches to improve species.

- Antonelli et al., 2016: Antonelli, L., Foata, J., Quilichini, Y., & March&, B. (2016). Influence of season & site location on European cultured sea bass parasites in Corsican fish farms using indicator species analysis (IndVal). *Parasite Research*, 115(2), 561-568. <https://doi.org/10.1007/s00436-015-4772-9>
- Bendtsen & Hansen, 2013: Bendtsen, J., & Hansen, J. L. S. (2013). A model of life cycle, connectivity & population stability of benthic macro-invertebrates in the North Sea/Baltic Sea transition zone. *Ecological Modelling*, 267, 54-65. <https://doi.org/10.1016/j.ecolmodel.2013.07.012>
- Bouchet, V. M. P., Telford, R. J., Rygg, B., Oug, E., & Alve, E. (2018). Can benthic foraminifera serve as proxies for changes in benthic macrofaunal community structure? Implications for the definition of reference conditions. *Marine Environmental Research*, 137, 24-36. <https://doi.org/10.1016/j.marenvres.2018.02.023>
- Burger, J. (2006). Bioindicators: Types, development, & use in ecological assessment & research. *Environmental Bioindicators*, 1(1), 22-39. <https://doi.org/10.1080/15555270590966483>
- Bustos-Baez, S., & Frid, C. (2003). Using indicator species to assess the state of macrobenthic communities: Advances in polychaete research. *Hydrobiologia*, 496(1-3), 299-309. <https://doi.org/10.1023/A:1026169520547>
- Carvalho, L. R. S., Loiola, M., & Barros, F. (2017). Manipulating habitat complexity to underst& its influence on benthic macrofauna. *Journal of Experimental Marine Biology & Ecology*, 489, 48-57. <https://doi.org/10.1016/j.jembe.2017.01.014>
- Castro, H., Ramalheira, F., Quintino, V., & Rodrigues, A. M. (2006). Amphipod acute & chronic sediment toxicity assessment in estuarine environmental monitoring: An example from Ria de Aveiro, NW Portugal. *Marine Pollution Bulletin*, 53(1-4), 91-99. [10.1016/j.marpolbul.2005.09.029](https://doi.org/10.1016/j.marpolbul.2005.09.029)
- Checon, H. H., Vieira, D. C., Corte, G. N., Sousa, E. C. P. M., Fonseca, G., & Amaral, A. C. Z. (2018). Defining soft bottom habitats & potential indicator species as tools for monitoring coastal systems: A case study in a subtropical bay. *Ocean & Coastal Management*, 164, 68-78. <https://doi.org/10.1016/j.ocecoaman.2018.03.035>
- Chen, L. L., Wang, Q. C., Li, X. J., Zhou, Z. Q., & Li, B. Q. (2016). Long-term trends of macrobenthos in southern Bohai Sea, China, in relation to environmental changes. *Scientia Sinica Vitae*, 46(9), 1121-1134. <https://doi.org/10.1360/N052016-00063>
- Borja, A., Franco, J., & Pérez, V. (2000). A marine biotic index to establish the ecological quality of soft-bottom benthos within European estuarine & coastal environments. *Marine pollution bulletin*, 40(12), 1100-1114. DOI: [10.1016/S0025-326X\(00\)00061-8](https://doi.org/10.1016/S0025-326X(00)00061-8)
- Chytrý, M., Tichý, L., Holt, J., & Botta-Dukát, Z. (2002). Determination of diagnostic species with statistical fidelity measures. *Journal of Vegetation Science*, 13(1), 79-90. <https://doi.org/10.1111/j.1654-1103.2002.tb02025.x>
- Coutellec, M., & Barata, C. (2013). Special issue on long-term ecotoxicological effects: An introduction. *Ecotoxicology*, 22(5), 763-766. <https://doi.org/10.1007/s10646-013-1092-7>
- De Cáceres, M., Legendre, P., & Moretti, M. (2010). Improving indicator species analysis by combining groups of sites. *Oikos*, 119(10), 1674-1684. <https://doi.org/10.1111/j.1600-0706.2010.18334.x>Digital Object Identifier (DOI)
- De la Vega, C., Paar, M., Köhler, L., von Dorrien, C., Kriegl, M., Oesterwind, D., & Schubert, H. (2023). Trophic redundancy in benthic fish food webs increases with scarcity of prey items, in the southern Baltic Sea. *Frontiers in Marine Science*, Article 1143792. <https://doi.org/10.3389/fmars.2023.1143792>
- Dean, H. K. (2015). The use of polychaetes (Annelida) as indicator species of marine pollution: A review. *Revista de Biología Tropical*, 56, 1-11. DOI: [10.15517/RBT.V56I4.27162](https://doi.org/10.15517/RBT.V56I4.27162)
- Dufrêne, M., & Legendre, P. (1997). Species assemblages & indicator species: The need for a flexible asymmetrical approach. *Ecological Monographs*, 67(3), 345-366. [https://doi.org/10.1890/0012-9615\(1997\)067\[0345:SAIST\]2.0.CO;2](https://doi.org/10.1890/0012-9615(1997)067[0345:SAIST]2.0.CO;2)
- Ellis, J., Anlauf, H., Kürten, S., Lozano-Cortés, D., Alsaffar, Z., Cúrdia, J., ... & Carvalho, S. (2017). Cross shelf benthic biodiversity patterns in the Southern Red Sea. *Scientific reports*, 7(1), 437. DOI: [10.1038/s41598-017-00507-y](https://doi.org/10.1038/s41598-017-00507-y)
- Fichet, D., Radenac, M.-H., & Miram&, P. (1998). Experimental studies of impacts of harbour sediments resuspension to marine invertebrates larvae: Bioavailability of Cd, Cu, Pb & Zn & toxicity. *Marine Pollution Bulletin*, 36(7), 509-518. [10.1016/S0025-326X\(97\)00190-2](https://doi.org/10.1016/S0025-326X(97)00190-2)

- Fonseca, G., & Gallucci, F. (2016). The need of hypothesis-driven designs & conceptual models in impact assessment studies: An example from the free-living marine nematodes. *Ecological Indicators*, 71, 79-86. <https://doi.org/10.1016/j.ecolind.2016.06.051>
- Gadelha, J. R., Morgado, F., & Soares, A. M. V. M. (2013). Histology & histochemistry of sea anemones in environmental contamination studies. *Microscopy & Microanalysis*, 19(S4), 57-58. <https://doi.org/10.1017/S1431927613000901>
- Gallo, L., Corapi, A., Loppi, S., & Lucadamo, L. (2014). Element concentrations in the lichen *Pseudevernia furfuracea* (L.) Zopf transplanted around a cement factory (Southern Italy). *Ecological Indicators*, 46, 566-574. <https://doi.org/10.1016/j.ecolind.2014.07.029>
- Geng, N., Wu, Y., Zhang, M., Tsang, D. C., Rinklebe, J., Xia, Y., ... & Ok, Y. S. (2019). Bioaccumulation of potentially toxic elements by submerged plants & biofilms: A critical review. *Environment international*, 131, 105015. DOI: 10.1016/j.envint.2019.105015
- Gérard, C., & Lance, E. (2019). Decline of freshwater gastropods exposed to recurrent interacting stressors implying cyanobacterial proliferations & droughts. *Aquatic Ecology*, 53(1), 79-96. <https://doi.org/10.1007/s10452-019-09674-8>
- Gesteira, J. G., & Dauvin, J. C. (2000). Amphipods are good bioindicators of the impact of oil spills on soft-bottom macrobenthic communities. *Marine Pollution Bulletin*, 40(11), 1017-1027. DOI: 10.1016/S0025-326X(00)00046-1
- Gislason, H., Bastardie, F., Dinesen, G. E., Egekvist, J., & Eigaard, O. R. (2017). Lost in translation? Multi-metric macrobenthos indicators & bottom trawling. *Ecological Indicators*, 82, 260-270. <https://doi.org/10.1016/j.ecolind.2017.07.004>
- Halpern, B. S., Frazier, M., Potapenko, J., Casey, K. S., Koenig, K., Longo, C., ... & Walbridge, S. (2015). Spatial & temporal changes in cumulative human impacts on the world's ocean. *Nature communications*, 6(1), 7615. <https://www.nature.com/articles/ncomms8615>
- Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Agrosa, C., Bruno, J. F., Casey, K. S., Ebert, C., Fox, H. E., Fujita, R., Heinemann, D., Lenihan, H. S., Madin, E. M. P., Perry, M. T., Selig, E. R., Spalding, M., Steneck, R., Watson, R. (2008). A global map of human impact on marine ecosystems. *Science*, 319(5865), 948-952. <https://doi.org/10.1126/science.1149345>
- Han, W., & Han, Q. (2024). Macrobenthic indicator species: From concept to practical applications in marine ecology. *Global Ecology & Conservation*, e03262.
- Han, W., Yan, R., Peng, S., Zhou, W., Yang, Y., & Han, Q. (2025). Integrated macrobenthic TITAN, AMBI, & M-AMBI indices for assessment of marine ecological quality in the East China Sea. *Marine Pollution Bulletin*, 214, 117704. 10.1016/j.marpolbul.2025.117704
- Hill, M. O. (1979). *Twinspan: A Fortran program for arranging multivariate data in an ordered two-way table by classification of individual & attributes*. Cornell University Press.
- ISO. (1979). International Organisation for Standardisation (ISO) (1979). Assessment of the biological quality of rivers by a macroinvertebrate "score". ISO/TC 147/SC 5/WG 6/N5 (Geneva: International Organisation for Standardisation, ISO) 18.
- Ito, M., Ito, K., Ohta, K., Hano, T., Onduka, T., Mochida, K., & Fujii, K. (2016). Evaluation of bioremediation potential of three benthic annelids in organically polluted marine sediment. *Chemosphere*, 163, 392-399. <https://doi.org/10.1016/j.chemosphere.2016.08.046>
- Keup, L. E., Mackenthun, K. M., & Ingram, W. M. (1967). *Biology of water pollution: A collection of selected papers on stream pollution, waste water, & water treatment*. United States Department of the Interior, Federal Water Pollution Control Administration.
- Khedhri, I., Atoui, A., Ibrahim, M., Afli, A., & Aleya, L. (2016). Assessment of surface sediment dynamics & response of benthic macrofauna assemblages in Boughrara Lagoon (SW Mediterranean Sea). *Ecological Indicators*, 70, 77-88.
- Kolkwitz, R., & Marsson, M. (1908). Ecological plant saprobionts. *Bericht der Deutschen Botanischen Gesellschaft*, 26a, 505-519.
- Kolkwitz, R., & Marsson, M. (1909). Ecology of animal saprobionts. *International Review of Hydrobiology & Hydrography*, 2, 126-152.
- L&res, P. B., Verner, J., & Thomas, J. W. (1988). Ecological uses of vertebrate indicator species: a critique. *Conservation biology*, 2(4), 316-328.

- Leonardsson, K., Blomqvist, M., & Rosenberg, R. (2009). Theoretical & practical aspects on benthic quality assessment according to the EU Water Framework Directive – Examples from Swedish waters. *Marine Pollution Bulletin*, 58(9), 1286-1296. <https://doi.org/10.1016/j.marpolbul.2009.05.007>
- Levin, L. A., & Gage, J. D. (1998). Relationships between oxygen, organic matter & the diversity of bathyal macrofauna. *Deep Sea Research Part II: Topical Studies in Oceanography*, 45(1-3), 129-163.
- Li, B., Jiang, S., Lü, J., Chen, L., Yan, L., Liu, C., Li, X., Song, B., & Li, X. (2020). Species composition & long-term variation of macrobenthos in intertidal zone & offshore areas of the Yellow River Delta. *Biodiversity Science*, 28(12), 1511–1522. <https://doi.org/10.17520/biods.2020164>
- Li, B., Li, X., Bouma, T. J., Soissons, L. M., Cozzoli, F., Wang, Q., Zhou, Z., & Chen, L. (2017). Analysis of macrobenthic assemblages & ecological health of Yellow River Delta, China, using AMBI & M-AMBI assessment method. *Marine Pollution Bulletin*, 119(2), 23–32. [10.1016/j.marpolbul.2017.03.044](https://doi.org/10.1016/j.marpolbul.2017.03.044)
- Li, B., Li, X., Bouma, T. J., Soissons, L. M., Cozzoli, F., Wang, Q., ... & Chen, L. (2017). Analysis of macrobenthic assemblages & ecological health of Yellow River Delta, China, using AMBI & M-AMBI assessment method. *Marine Pollution Bulletin*, 119(2), 23-32.
- Li, B., Wang, Q., & Li, B. (2013). Assessing the benthic ecological status in the stressed coastal waters of Yantai, Yellow Sea, using AMBI & M-AMBI. *Marine pollution bulletin*, 75(1-2), 53-61.
- Li, F., Ma, Y., Song, X., Li, S., Zhang, X., Wang, X., Wang, T., & Sun, Z. (2022). Community structure & ecological quality assessment of macrobenthos in the coastal sea areas of Northern Yantai, China. *Frontiers in Marine Science*, 9, 989034. <https://doi.org/10.3389/fmars.2022.989034>
- Liu, C., Xu, Q., Wang, Z., Jiang, X., Ding, G., Ren, Q., Song, J., & Liu, M. (2023). Community structure of benthic molluscs shaped by environmental & ecological variables in the coastal waters of Changle, Fujian Province, China. *Frontiers in Marine Science*, 10, Article 1045393. <https://doi.org/10.3389/fmars.2023.1045393>
- Liu, L. S., Li, Z. Y., Meng, W., Zheng, B. H., & Hu, X. A. (2007). The community structure of zoobenthos & bioassessment of water quality in the lower reaches of the Songhua River. *Research of Environmental Sciences*, 20(3), 81-86.
- Liu, L. S., Meng, W., Li, X. Z., Li, Z. C., Zheng, B. H., Lei, K., & Li, Z. Y. (2009). Studies on macrobenthos in the northern waters of Liaodong Bay: II. Biodiversity & community structure. *Research of Environmental Sciences*, 22(2), 155-161.
- Liu, X. D., Wang, J. S., Sun, L. E., Cui, W. L., & Li, Z. (2021). Community structure & diversity of macrozoobenthos at Bohai Sea off Sh&ong. *Marine Environmental Science*, 40(6), 929-936.
- Liu, X. S., Zhao, R., Hua, E., Lu, L., & Zhang, Z. N. (2014). Macrofaunal community structure in the Laizhou Bay in summer & the comparison with historical data. *Mar. Sci. Bull*, 33(10).
- Luo, X. X., & Yang, J. Q. (2009). Progress in researches on benthic indices of assessing marine ecosystem health. *Marine Science Bulletin*, 28(3), 106-112.
- Maanan, M. (2007). Biomonitoring of heavy metals using *Mytilus galloprovincialis* in Safi coastal waters, Morocco. *Environmental Toxicology*, 22(5), 525–531. <https://doi.org/10.1002/tox.20301>
- Martin, J. P., Lizarralde, Z. I., Pittaluga, S., Sar, A. M., Perroni, M., & Torres, S. (2019). Macrobenthic assemblages of intertidal flats in Southern Patagonia, Argentina. *Thalassas: An International Journal of Marine Sciences*, 35, 689-705.
- Marujo Neves, J., Martins, M., Adão, H., Mil-Homens, M., Costa, M. H., & Lobo-Arteaga, J. (2025). Ecological quality assessment of estuarine macrobenthic communities using an integrative approach. *Marine Pollution Bulletin*, 210, 117316.
- McArthur, M. A., Brooke, B. P., Przeslawski, R., Ryan, D. A., Lucieer, V. L., Nichol, S., McCallum, A. W., Mellin, C., Cresswell, I. D., Radke, L. C. (2010). On the use of abiotic surrogates to describe marine benthic biodiversity. *Estuarine, Coastal & Shelf Science*, 88(1), 21-32. <https://doi.org/10.1016/j.ecss.2010.03.003>
- Nasi, F., Nordström, M. C., Bonsdorff, E., Auriemma, R., Cibic, T., & Del Negro, P. (2018). Functional biodiversity of marine soft-sediment polychaetes from two Mediterranean coastal areas in relation to environmental stress. *Marine Environmental Research*, 137, 121-132. <https://doi.org/10.1016/j.marenvres.2018.03.002>
- Podlesińska, W., & Dąbrowska, H. (2019). Amphipods in estuarine & marine quality assessment – a review. *Oceanologia*, 61(2), 179-196. <https://doi.org/10.1016/j.oceano.2018.09.002>

- Radenac, G.-H., Fichet, D., & Miramand, P. (2001). Bioaccumulation & toxicity of four dissolved metals in *Paracentrotus lividus* sea-urchin embryo. *Marine Environmental Research*, 51(2), 151–166. 10.1016/S0141-1136(00)00092-1
- Rakaj, A., Morroni, L., Grosso, L., Fianchini, A., Pensa, D., Pellegrini, D., & Regoli, F. (2021). Towards sea cucumbers as a new model in embryo-larval bioassays: *Holothuria tubulosa* as test species for the assessment of marine pollution. *Science of the Total Environment*, 787, Article 147593. <https://doi.org/10.1016/j.scitotenv.2021.147593>
- Reish, D. I. (1985). A discussion of the importance of screen size in washing quantitative marine bottom samples. *Ecology*, 46(2), 307–309. <https://doi.org/10.2307/1930047>
- Somers, H. L. (1958). Benthic Studies in Buzzards Bay. I. Animal-Sediment Relationships 1. *Limnology & oceanography*, 3(3), 245–258.
- Siddig, A. A. H., Ellison, A. M., Ochs, A., Villar-Leeman, C., & Lau, M. K. (2016). How do ecologists select & use indicator species to monitor ecological change? Insights from 14 years of publication in ecological indicators. *Ecological Indicators*, 60, 223–230. <https://doi.org/10.1016/j.ecolind.2015.06.036>
- Sobczyk, R., Czortek, P., Serigstad, B., & Pabis, K. (2021). Modelling of polychaete functional diversity: Large marine ecosystem response to multiple natural factors & human impacts on the West African continental margin. *Science of the Total Environment*, 792, Article 148075. <https://doi.org/10.1016/j.scitotenv.2021.148075>
- Sukumaran, S., Vijapure, T., Mulik, J., & Ridha, H. (2021). Marine macrobenthos of North West India- Reviewing the known & unknown. *Frontiers in Marine Science*, 8, 671245.
- Szefer, P., Kim, B.-S., Kim, C.-K., Kim, E.-H., & Lee, C.-B. (2005). Biomonitoring of trace metals in the Black Sea (Romania) using mussels *Mytilus galloprovincialis*. *Water Research*, 39(4), 596–604. <https://doi.org/10.1016/j.watres.2004.09.026>
- Wang, Z., Wang, H., Fan, S., Xin, M., & Sun, X. (2021). Community structure & diversity of macrobenthos in Jiaozhou Bay. *Marine Pollution Bulletin*, 171, 112781.
- Wong, C. K. C., Cheung, R. Y. H., & Wong, M. H. (2005). Biomonitoring of heavy metals & trace organics using the intertidal mussel *Perna viridis* in Hong Kong coastal waters. *Marine Pollution Bulletin*, 51(8–12), 857–875. 10.1016/j.marpolbul.2005.04.014
- Xu, Y., & Li, X. Z. (2021). Macrobenthos & offshore ecosystem health evaluation. *Science*, 73, 30–34.
- Yan, J., Sui, J., Xu, Y., Li, X., Wang, H., & Zhang, B. (2020). Assessment of the benthic ecological status in adjacent areas of the Yangtze River Estuary, China, using AMBI, M-AMBI & BOPA biotic indices. *Marine Pollution Bulletin*, 157, 111020. 10.1016/j.marpolbul.2020.111020
- Yang, C., Ding, S., Niu, Y., Zhang, X., & Li, J. (2024). Characteristics of Macrobenthic Communities & their Response to Environmental Changes in East Taihu Lake, China.
- Yap, C. K., Ismail, A., & Tan, S. G. (2003). Can the byssus of green-lipped mussel *Perna viridis* (Linnaeus) from the west coast of Peninsular Malaysia be a biomonitoring organ for Cd, Pb & Zn? Field & laboratory studies. *Environment International*, 29(4), 521–528. 10.1016/S0160-4120(03)00008-4
- Zhang, W., Liu, X., Cheng, H., Zang, J., & Xu, H. (2012). Distribution & community structure of macrobenthos in relation to environmental factors in the Yellow Sea. *Acta Ecologica Sinica*, 32, 4050–4061
- Zhang, Y., Li, S. W., Lv, Z. B., Ma, Y. Q., Liu, Y. J., Wei, Z. H., & Chen, W. (2013). Application of polychaete in ecological environment evaluation of Laizhou Bay. *Acta Ecol. Sin.*, 33, 2522–2530.